DEPTH DISCRIMINATION AND RANGE ESTIMATION OF MULTIPLE SOURCES IN SHALLOW WATERS

I-Tai Lu

LONG-TERM GOALS

To develop wave-oriented phase coherent signal processing algorithms (which address and incorporate the physics of acoustic wavefield propagation through and scattering from an inhomogeneous, noisy, time varying, and stochastic ocean environment) to characterize targets and the environment for applications in both passive and active systems.

OBJECTIVES

To validate a new and practical system concept and to develop new wave-oriented phase coherent inversion algorithms for classifying and localizing multiple sources in noisy and time-varying shallow waters without knowing the sound speed profiles. The approaches should be able to adapt (self-optimize) in the face of changing environmental and system conditions.

APPROACHES

There have been many efforts to develop inverse source algorithms such as matched-field, matched-traveltime, matched-ray, and matched-mode approaches. However, none of the existing methods can work well in real time under simultaneous excitations of several sources. Furthermore, all of these algorithms operate properly only when accurate descriptions of the environments are available. Since the environment is time-varying and full of noises created by surface ships and other sources, it is impractical to implement any real time monitoring system using these methods. A real time monitoring system with two mutually perpendicular horizontal arrays (which can be bottom mounted) is proposed for range estimation, depth discrimination, and spectrum estimation of multiple sources where the SSP of the environment is not required. The key ideas are, first, to use sufficient long arrays for deriving modal wave numbers and composite modal amplitudes excited by multiple sources in multiple frequencies and, then, to utilize a sufficient large number of frequency samples for obtaining unknown source ranges and modal amplitudes for every source and every guided mode.

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number	ion of information Send comment arters Services, Directorate for Inf	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	his collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE 30 SEP 1997	2 DEPORT TYPE			3. DATES COVERED 00-00-1997 to 00-00-1997		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
Depth Discrimination and Range Estimation of Multiple Sources in Shallow Waters				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Polytechnic University,Route 110,Farmingdale,NY,11735				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release; distributi	ion unlimited				
13. SUPPLEMENTARY NO	OTES					
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON		
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified	Same as Report (SAR)	3	REST ONSIBLE I ERSON	

Report Documentation Page

Form Approved OMB No. 0704-0188

WORK COMPLETED

The proposed system is tested numerically in an simplified shallow water model. The conventional spectral estimation model is extended to incorporate the array coordinate dependence for wavenumber inversion and to account for the frequency dependence for range estimation. A new Singular Value Decomposition - Eigenmatrix Pencil method is employed to find the complex modal wavenumber and the complex source ranges. This high-resolution algorithm is proven to be most successful for the current application.

RESULTS

In numerical simulations, the power spectrum and the range for each source are approximately derived from its fundamental modal phase and amplitude, respectively. Furthermore, depth discrimination is achieved by examining the excitation strength of the fundamental mode especially when the environment has a downward refracting SSP. If the SSP or the modal eigenfunctions is approximately known, more accurate source depths can be derived from modal excitation strengths by a matched-mode procedure.

IMPACT/APPLICATIONS

Source localization technologies are of vital importance to the US Navy. The success of the preliminary results of the proposed research program is a fundamental breakthrough in inverse source algorithms. These results will enable the US Navy to develop inexpensive real time systems to monitor broad regions in crucial and busy shallow waters so as to detect underwater vessels, estimate their ranges and acoustic power spectra, and discriminate them from surface ships. These results may also be employed for tracking whales or for other civilian applications.

REFERENCES

A. Journal Papers:

- 1. <u>I.T. Lu</u> and Y. Q. Yao, "Ray shooting, eigenray search, and complex resonances of submerged structures of revolution," J. Acoust. Soc. Am, 101(1), 66-76, 1997.
- 2. <u>I.T. Lu</u>, "Simultaneous characterization of source and environment using a ray travel-time inversion approach," J. Computational Acoustics, Vol. 5, No. 2, 193-218, 1997.
- 3. C. M. Qiu and <u>I. T. Lu</u>, "Path frequency consideration for the RAKE receiver in a CDMA cellular system," in press, IEEE Trans. Veh. Techno.
- 4. <u>I. T. Lu</u> and C. M. Qiu, "A novel high-resolution algorithm for complex-direction search," submitted to J. Acoust. Soc. Am.

B. Conference Papers:

- 1. <u>I. T. Lu</u> and C. M. Qiu, "A novel high-resolution algorithm for complex-direction serach," 1996 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP.96), Atlanta, GA.
- 2. <u>I-Tai Lu</u> and Hsuan-Ling Wu, Range Estimation of Multiple Sources in Shallow Waters Without Knowing the Sound Speed Profile, 'Third International Conference on Theoretical & Computational Acoustics, Columbia Univ., New York, July 14-18, 1997.
- 3. <u>I-Tai Lu</u> and Fu-Po Wu, "A High-Resolution Spectral Estimation Approach for Evaluating the Traveltimes and Resonances of Scattering Returns from Submerged Targets," Third International Conference on Theoretical & Computational Acoustics, Columbia Univ., New York, July 14-18, 1997.
- 4. <u>I-Tai Lu</u> and T. C. Yang, "Alternative Representations of Acoustics Field Scattered by Submerged Targets in Ocean Waveguides," Third International Conference on Theoretical & Computational Acoustics, Columbia Univ., New York, July 14-18, 1997.